

Technical Procedure for Coomassie Blue

1.0 Purpose - This procedure describes how to make Coomassie Blue solution and apply it to items of evidence.

2.0 Scope – This procedure applies to porous and non-porous items of evidence that may contain bloody impressions that require developing/enhancing. This procedure may also be used in processing adhesive sides of tapes.

3.0 Definitions – N/A

4.0 Equipment, Materials and Reagents

4.1 Equipment and Materials

- Protective gloves and apron/coat
- Face shield visor and/or safety goggles
- Magnetic stirrer, magnetic follower and magnetic retriever
- Two (2) glass beakers
- Application equipment: two (2) spray bottles and two (2) glass trays
- Camera/scanner
- Fume hood

4.2 Reagents

- Coomassie Brilliant Blue (0.44 gram)
- Glacial acetic acid (40 mL)
- Methanol (200 mL)
- Distilled water (200 mL)

5.0 Procedure

5.1 Mixing Procedure

5.1.1 Staining Solution (Developer)

5.1.1.1 Place 0.44 g of Coomassie Brilliant Blue and two-hundred (200) mL of methanol in a large glass beaker with magnetic stirrer and stir.

5.1.1.2 Add forty (40) ml of glacial acetic acid and two-hundred (200) mL of distilled water to the solution with continuous stirring to ensure the solution is thoroughly mixed.

5.1.1.3 Place the solution in a clearly marked spray bottle for immediate use or a dark container for long term use as needed.

5.1.2 Destaining Solution (Rinse Solution)

5.1.2.1 Place forty (40) mL of glacial acetic acid and two-hundred (200) mL of methanol in a large beaker with a magnetic stirrer and stir.

- 5.1.2.2 Add two-hundred (200) mL of distilled water to the solution with continuous stirring to ensure the solution is thoroughly mixed.
- 5.1.2.3 Place the solution in a clearly marked spray bottle for immediate use or a dark jug for long term use as needed.
- 5.1.2.4 Larger amounts of Coomassie Blue may be mixed for large items or for use at crime scenes.

5.2 Application Procedure Forensic Scientists shall produce a self-made test print to be processed concurrently with items of evidence. (See Section Technical Procedure for Ensuring Quality Control.)

5.2.1 Spray Method

- 5.2.1.1 Completely cover the area of interest with the staining solution.
- 5.2.1.2 Spray the item with the destaining solution to clear the background. The destaining solution shall be used generously to remove the excess staining solution.
- 5.2.1.3 Allow the item to dry completely prior to proceeding.

5.2.2 Immersion Method

- 5.2.2.1 Completely immerse the item in a tray of staining solution. Immerse the item for approximately thirty (30) to ninety (90) seconds.
- 5.2.2.2 Remove the item from the staining solution and place in a separate tray of destaining solution for approximately one minute and agitate to clear the background. This procedure may be repeated with a fresh destaining solution if the background is not completely clear.
- 5.2.2.3 Remove the item from the solution and allow the item to dry completely prior to proceeding.
- 5.2.2.4 Preserve the developed impressions through photography, according to the techniques in Photographic Equipment/Procedures and/or by electronically recording the impressions (See Image Processing). The impression may be lifted directly from the item only after the item is completely dry.

5.3 Standards and Controls – N/A

5.4 Calibration - N/A

5.5 Sampling –N/A

5.6 Calculations – N/A

5.7 Uncertainty of Measurement – N/A

6.0 Limitations - Coomassie Blue powder and solution have an indefinite shelf life.

6.1 The Coomassie Blue reagent shall be stored in the original shipping container until needed.

6.2 The staining and destaining solutions may be stored in clear spray bottles or dark containers until needed.

7.0 Safety – Glacial acetic acid and methanol can be harmful if inhaled or ingested and shall be used in a fume hood when mixing and/or processing evidence. Protective gloves, eye goggles and aprons shall be worn as the staining solution will stain clothing and skin. This technique may be used at crime scenes; however, use only in a well vented area or use a fan to remove the fumes produced.

8.0 References

Hunter, J.L. “Fingerprint Evidence with Coomassie Blue—After 25 years.” *Journal of Forensic Identification*. Vol. 44, 6: 619-622 (1994).

Kent, T. ed. *Manual of Fingerprint Development Techniques: A Guide to the Selection and Use of Processing for the Development of Latent Fingerprints*. Police Scientific Development Branch, London (July 1992).

Lee, H.C. “Methods of Latent Print Development.” *Proceedings of the International Forensic Symposium on Latent Prints*. (July 1987): 15 – 24.

Lennard, C.J. and P.A. Margot. “Sequencing of Reagents for the Improved Visualization of Latent Fingerprints.” *Proceedings of the International Forensic Symposium on Latent Prints*. (July 1987): 141-142.

Manual of Fingerprint Development Techniques. (January 1986): 2-8.

Manual of Fingerprint Development Techniques: A Guide to the Selection and Use of Processes for the Development of Latent Fingerprints. Scientific Research and Development Branch, London (1986).

Norkus, P. M. and K. Noppinger. “Enhanced Latent Prints in Blood with a New Staining Technique.” *Proceedings of the International Forensic Symposium on Latent Prints*. (1987): 147.

Trozzi, T. A., R.L. Schwartz and M.L. Hollars. *Processing Guide for Developing Latent Prints*. (2000): 1-64.

US Department of Justice. *Chemical Formulas and Processing Guide for Developing Latent Prints*. FBI Laboratory Division, Latent Fingerprint Section (1994).

9.0 Records – N/A

10.0 Attachment – N/A

Revision History		
Effective Date	Version Number	Reason
09/17/2012	1	Original Document
10/31/2013	2	Added issuing authority to header